



TITLE:

# On certain subclasses of analytic functions involving a linear operator

AUTHOR(S):

Saitoh, Hitoshi; Nunokawa, Mamoru

---

CITATION:

Saitoh, Hitoshi ...[et al]. On certain subclasses of analytic functions involving a linear operator. 数理解析研究所講究録 1996, 963: 97-109

ISSUE DATE:

1996-08

URL:

<http://hdl.handle.net/2433/60555>

RIGHT:

## On certain subclasses of analytic functions involving a linear operator

群馬高専

斎藤 斉 (Hitoshi Saitoh)

群馬大・教育

布川 護 (Mamoru Nunokawa)

### Abstract

A certain linear operator defined by a Hadamard product or convolution for functions which are analytic in the open unit disk is introduced according to Carlson and Shaffer. The purpose of the present paper is to give some properties of this linear operator. Our results contain the earlier theorems in the univalent and multivalent function theory.

# 1 Introduction

Let  $A_p$  denote the class of functions of the form

$$f(z) = z^p + \sum_{n=1}^{\infty} a_{n+p} z^{n+p} \quad (p \in N = \{1, 2, 3, \dots\}) \quad (1.1)$$

which are analytic in the unit disc  $U = \{z : |z| < 1\}$ , and  $A_1 = A$ .

For

$$f(z) = \sum_{n=0}^{\infty} a_n z^n \quad \text{and} \quad g(z) = \sum_{n=0}^{\infty} b_n z^n ,$$

we define the Hadamard product ( or convolution ) by

$$(f * g)(z) = \sum_{n=0}^{\infty} a_n b_n z^n .$$

Let

$$\phi_p(a, c, ; z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} z^{n+p} \quad (c \neq 0, -1, -2, \dots; \quad z \in U) , \quad (1.2)$$

where  $(x)_n$  is the Pochhammer symbol defined by

$$(x)_n = \begin{cases} 1 & (\text{if } n = 0) \\ x(x+1)(x+2) \cdots (x+n-1) & (\text{if } n \in N) \end{cases} .$$

We note that

$$\phi_p(a, c; z) = z^p \cdot {}_2F_1(1, a; c; z) ,$$

where

$${}_2F_1(1, a; c; z) = \sum_{n=0}^{\infty} \frac{(1)_n (a)_n}{(c)_n} \frac{z^n}{n!} .$$

**Remark 1.**

$$\begin{aligned} \phi_p(a, 1; z) &= \frac{z^p}{(1-z)^a} \\ \phi_p(a, 1; z) &= \frac{z}{(1-z)^2} \quad (\text{koebe function}) . \end{aligned}$$

Corresponding to the function  $\phi_p(a, c; z)$ , we define a linear operator  $L_p(a, c)$  on  $A_p$  by the convolution

$$L_p(a, c)f(z) = \phi_p(a, c; z) * f(z) \quad (1.3)$$

for  $f(z) \in A_p$ .

**Remark 2.** If  $c > a > 0$ ,  $L_p(a, c)$  has integral representation

$$L_p(a, c)f(z) = \int_0^1 u^{-p} f(uz) d\mu(u),$$

where  $\mu$  satisfies

$$d\mu(u) = \frac{u^{a-1}(1-u)^{c-a-1}}{B(a, c-a)} du \quad \text{and} \quad \int_0^1 d\mu(u) = 1.$$

Clearly,  $L_p(a, a)$  is the unit operator and

$$L_p(a, c) = L_p(a, b)L_p(b, c) = L_p(b, c)L_p(a, b) \quad (b, c \neq 0, -1, -2, \dots).$$

Moreover, if  $a \neq 0, -1, -2, \dots$ , then  $L_p(a, c)$  has an inverse  $L_p(c, a)$ .

The operator  $L_1(a, c)$  was introduced by Carlson and Shaffer [2] in their systematic investigation of certain interesting classes of starlike, convex and prestarlike hypergeometric functions.

In recent years Srivastava and Owa [15] have given some properties of  $L_1(a, c)$  concerning with univalent functions in  $U$ .

**Remark 3.** For  $f(z) \in A_p$ ,

$$L_p(\nu + p, 1)f(z) = \frac{z^p}{(1-z)^{\nu+p}} * f(z) = D^{\nu+p-1}f(z), \quad (1.4)$$

where  $\nu(>p)$  is any real number. In the case of  $p = 1$  and  $\nu \in N$ ,  $D^\nu f(z)$  is the Ruscheweyh derivative [8].

$$L_p(\nu + p, \nu + p + 1)f(z) = \frac{\nu + p}{z^\nu} \int_0^z t^{\nu-1} f(t) dt = J_{\nu,p}(f(z)), \quad (1.5)$$

where  $\nu + p > 0$ . The operator  $J_{\nu,1}(\nu \in N)$  was introduced by Bernardi [1]. In particular, the operator  $J_{1,1}$  was studied earlier by Libera [4] and Livingston [5]. Some results for the operator  $J_{\nu,p}$  were showed by Saitoh [10] and Saitoh et al [14].

A function  $f(z)$  belonging to the class  $A_1$  is said to be starlike if and only if it satisfies

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0 \quad (z \in U). \quad (1.6)$$

A function  $f(z)$  belonging to the class  $A_1$  is convex if and only if it satisfies

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > 0 \quad (z \in U). \quad (1.7)$$

The subclass of univalent functions consisting of the starlike functions is denoted by  $S^*$ , and  $K$  denotes the subclass of convex functions.

A function  $f(z) \in A_p$  is said to be  $p$ -valently starlike if and only if it satisfies

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0 \quad (z \in U) . \quad (1.8)$$

A function  $f(z) \in A_p$  is convex if and only if it satisfies

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > 0 \quad (z \in U) . \quad (1.9)$$

## 2 Some properties of univalent and multivalent functions involving the operator $L_p(a, c)$

In order to prove our results we need the following lemmas.

**Lemma 1.** (Saitoh [11]) *If  $f(z) \in A_p$ , then*

$$z(L_p(a, c)f(z))' = aL_p(a+1, c)f(z) - (a-p)L_p(a, c)f(z) \quad (2.1)$$

where  $c \neq 0, -1, -2, \dots$ .

**Proof.** Note that

$$L_p(a, c)f(z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} a_{n+p} z^{n+p} \quad (a_p = 1)$$

and

$$L_p(a+1, c)f(z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} a_{n+p} z^{n+p} \quad (a_p = 1) .$$

These give that

$$\begin{aligned} & aL_p(a+1, c)f(z) - (a-p)L_p(a, c)f(z) \\ &= \sum_{n=0}^{\infty} (a+n) \frac{(a)_n}{(c)_n} a_{n+p} z^{n+p} - \sum_{n=0}^{\infty} (a-p) \frac{(a)_n}{(c)_n} a_{n+p} z^{n+p} \\ &= \sum_{n=0}^{\infty} (n+p) \frac{(a)_n}{(c)_n} a_{n+p} z^{n+p} \\ &= z(L_p(a, c)f(z))' . \end{aligned}$$

**Lemma 2.** (Saitoh and Nunokawa [12]) *Let  $g(z)$  be analytic and satisfies  $\operatorname{Re}\{g(z)\} > 0$  in  $U$ .*

*Then we have*

$$|zg'(z)| \leq \frac{2|z|\operatorname{Re}\{g(z)\}}{1-|z|^2} \quad (z \in U) . \quad (2.2)$$

**Lemma 3.** If  $g(z) = 1 + b_1z + b_2z^2 + \dots$  is analytic in  $U$  and  $\operatorname{Re}\{g(z)\} > 0$ . Then

$$\operatorname{Re}\{g(z)\} \geq \frac{1 - |z|}{1 + |z|} . \quad (2.3)$$

**Lemma 4.** (MacGregor [7]) Suppose that  $h(z) = 1 + c_1z + c_2z^2 + \dots$  is analytic and satisfies  $\operatorname{Re}\{g(z)\} > 0$  in  $U$ . Then

$$\left| \frac{h'(z)}{h(z)} \right| \leq \frac{2}{1 - |z|^2} \quad (z \in U) . \quad (2.4)$$

Applying lemmas, we prove

**Theorem 1.** Let  $f(z) \in A_p$  and  $a > 0$ . If

$$\operatorname{Re} \left\{ \frac{L_p(a, c)f(z)}{z^p} \right\} > 0 \quad (z \in U) ,$$

then

$$\begin{aligned} \operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{z^p} \right\} &\geq \frac{a - 2r - ar^2}{a(1+r)^2} \\ \text{for } |z| = r &< \frac{\sqrt{1+a^2} - 1}{a} . \end{aligned} \quad (2.5)$$

**Proof.** We put  $g(z) = \frac{L_p(a, c)f(z)}{z^p}$ .

Then  $g(z) = 1 + b_1z + b_2z^2 + \dots$  is analytic and satisfies  $\operatorname{Re}\{g(z)\} > 0$  in  $U$ .

Differentiating of  $z^p g(z) = L_p(a, c)f(z)$  and using Lemma 1, we have

$$\frac{L_p(a+1, c)f(z)}{z^p} = g(z) + \frac{1}{a}zg'(z) .$$

Therefore, using Lemma 2 and Lemma 3

$$\begin{aligned} \operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{z^p} \right\} &\geq \operatorname{Re}\{g(z)\} - \frac{1}{a}|zg'(z)| \\ &\geq \operatorname{Re}\{g(z)\} - \frac{2|z|\operatorname{Re}\{g(z)\}}{a(1-|z|^2)} \\ &= \frac{\operatorname{Re}\{g(z)\}(a - 2|z| - a|z|^2)}{a(1-|z|^2)} \\ &\geq \frac{1-|z|}{1+|z|} \frac{a - 2|z| - a|z|^2}{a(1-|z|^2)} \\ &= \frac{a - 2r - ar^2}{a(1+r)^2} \\ \text{for } |z| = r &< \frac{\sqrt{1+a^2} - 1}{a} . \end{aligned}$$

This completes the proof of Thm. 1.

Putting  $a = c = p$  in Thm. 1, we have

**Corollary 1.** (Saitoh and Nunokawa [12]) Let  $f(z) \in \mathbf{A}_p$ . If

$$\operatorname{Re} \left\{ \frac{f(z)}{z^p} \right\} > 0 \quad (z \in U) ,$$

then we have

$$\operatorname{Re} \left\{ \frac{f'(z)}{z^{p-1}} \right\} \geq \frac{p - 2r - pr^2}{(1+r)^2} \quad \text{for } |z| = r < \frac{\sqrt{1+p^2} - 1}{p} . \quad (2.6)$$

**Corollary 2.** (Yamaguchi [16]) Let  $f(z) = z + a_2 z^2 + \dots$  be analytic in  $U$ . If  $\operatorname{Re} \left\{ \frac{f(z)}{z} \right\} > 0$ , then we have

$$\operatorname{Re} \{f'(z)\} > 0 \quad \text{for } |z| = r < \sqrt{2} - 1 .$$

Next, we prove

**Theorem 2.** Let  $f(z) \in \mathbf{A}_p$  and  $a > 0$ . If

$$\operatorname{Re} \left\{ \frac{L_p(a, c)f(z)}{z^p} \right\} > 0 \quad (z \in U) ,$$

then

$$\left| \frac{aL_p(a+1, c)f(z)}{L_p(a, c)f(z)} - a \right| \leq \frac{2|z|}{1-|z|^2} \quad (z \in U) . \quad (2.7)$$

**Proof.** We put  $h(z) = \frac{L_p(a, c)f(z)}{z^p}$ . Then  $h(z) = 1 + c_1 z + c_2 z^2 + \dots$  is analytic and satisfies  $\operatorname{Re} \{h(z)\} > 0$  in  $U$ . Logarithmically differentiating of  $h(z)$  and using Lemma 1, we can see

$$\frac{aL_p(a+1, c)f(z)}{L_p(a, c)f(z)} - a = \frac{zh'(z)}{h(z)} .$$

From Lemma 4, we have

$$\left| \frac{aL_p(a+1, c)f(z)}{L_p(a, c)f(z)} - a \right| = \left| \frac{zh'(z)}{h(z)} \right| \leq \frac{2|z|}{1-|z|^2} .$$

This completes the proof of Thm. 2.

Letting  $a = c = p$  in Thm. 2, we have the following interesting Corollary.

Corollary 3. Let  $f(z) \in A_p$ . If

$$\operatorname{Re} \left\{ \frac{f(z)}{z^p} \right\} > 0 \quad (z \in U) ,$$

then  $f(z)$  is  $p$ -valently starlike in  $|z| < \frac{\sqrt{1+p^2}-1}{p}$ .

Proof. From Theorem 2,

$$\left| \frac{zf'(z)}{f(z)} - p \right| \leq \frac{2|z|}{1-|z|^2} < p \text{ for } |z| < \frac{\sqrt{1+p^2}-1}{p}$$

i.e.,  $\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0$  in  $|z| < \frac{\sqrt{1+p^2}-1}{p}$ .

Further, taking the function  $f(z)$  given by

$$f(z) = \frac{z^p(1-z)}{1+z} ,$$

we see that the result is sharp.

Corollary 4. (MacGregor [6]) Let  $f(z) = z + a_2z^2 + \dots$  be analytic and satisfies

$$\operatorname{Re} \left\{ \frac{f(z)}{z} \right\} > 0 \text{ in } U. \text{ Then } f(z) \text{ is starlike in } |z| < \sqrt{2}-1.$$

Using the same technique of Thm. 2, we prove

Theorem 3. Let  $f(z) \in A_p, a > 0$ . If

$$\operatorname{Re} \left\{ \frac{L_p(a, c)f(z)}{z^p} \right\} > 0 \quad (z \in U) ,$$

then

$$\operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{L_p(a, c)f(z)} \right\} \geq \frac{a-2|z|-a|z|^2}{a(1-|z|^2)} \quad (z \in U) . \quad (2.8)$$

Proof. We put  $h(z) = \frac{L_p(a, c)f(z)}{z^p}$ .

Then  $h(z) = 1 + c_1z + c_2z^2 + \dots$  is analytic and satisfies  $\operatorname{Re} \{h(z)\} > 0$  in  $U$ .

Differentiating of  $h(z)$  and using Lemma 1, we have

$$\frac{L_p(a+1, c)f(z)}{L_p(a, c)f(z)} = 1 + \frac{1}{a} \frac{zh'(z)}{h(z)} .$$

Therefore, from Lemma 4 we can see that

$$\begin{aligned} \operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{L_p(a, c)f(z)} \right\} &\geq 1 - \frac{1}{a} \left| \frac{zh'(z)}{h(z)} \right| \\ &\geq 1 - \frac{1}{a} \frac{2|z|}{1-|z|^2} \\ &= \frac{a-2|z|-a|z|^2}{a(1-|z|^2)} \end{aligned}$$



This completes the proof of Thm. 3.

Making  $a = c = p$  in Thm. 3, we can show Cor. 3.

Next, we prove

**Theorem 4.** Let  $f(z), g(z) \in A_p, a > 0$  and satisfies

$$\operatorname{Re} \left\{ \frac{L_p(a+1, c)g(z)}{L_p(a, c)g(z)} \right\} > 0 \quad (z \in U) .$$

If

$$\operatorname{Re} \left\{ \frac{L_p(a, c)f(z)}{L_p(a, c)g(z)} \right\} > 0 \quad (z \in U) ,$$

then

$$\operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{L_p(a, c)f(z)} \right\} > 0 \quad \text{for } |z| < \frac{a+1-\sqrt{2a+1}}{a} . \quad (2.9)$$

**Proof.** We put  $p(z) = \frac{L_p(a, c)f(z)}{L_p(a, c)g(z)}$ . Then  $p(z) = 1 + p_1z + p_2z^2 + \dots$  is analytic and satisfies  $\operatorname{Re}\{p(z)\} > 0$  in  $U$ . Differentiating of

$$L_p(a, c)f(z) = p(z) \cdot L_p(a, c)g(z) ,$$

we have

$$z(L_p(a, c)f(z))' = zp'(z)L_p(a, c)g(z) + p(z)z(L_p(a, c)g(z))'$$

From Lemma 1, we can see

$$\frac{L_p(a+1, c)f(z)}{L_p(a, c)f(z)} = \frac{L_p(a+1, c)g(z)}{L_p(a, c)g(z)} + \frac{1}{a} \frac{zp'(z)}{p(z)} .$$

Now, we put  $q(z) = \frac{L_p(a+1, c)g(z)}{L_p(a, c)g(z)}$ .

Then  $q(z) = 1 + b_1z + b_2z^2 + \dots$  is analytic and satisfies  $\operatorname{Re}\{q(z)\} > 0$  in  $U$ .

Therefore, using Lemma 3 and Lemma 4, we have

$$\begin{aligned} \operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{L_p(a, c)f(z)} \right\} &\geq \operatorname{Re}\{q(z)\} - \frac{1}{a} \left| \frac{zp'(z)}{p(z)} \right| \\ &\geq \frac{1-|z|}{1+|z|} - \frac{1}{a} \frac{2|z|}{1-|z|^2} \\ &= \frac{a|z|^2 - 2(a+1)|z| + a}{a(1-|z|^2)} \\ &> 0 \quad \text{in } |z| < \frac{a+1-\sqrt{2a+1}}{a} . \end{aligned}$$

This completes the proof of Thm. 4.

Putting  $a = c = p$  in Thm. 4, we have

**Corollary 5.** Let  $f(z), g(z) \in A_p$  and satisfies

$$\operatorname{Re} \left\{ \frac{zg'(z)}{g(z)} \right\} > 0 \quad (z \in U) .$$

If

$$\operatorname{Re} \left\{ \frac{f(z)}{g(z)} \right\} > 0 \quad (z \in U) ,$$

then

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0 \text{ in } |z| < \frac{p+1-\sqrt{2p+1}}{p} . \quad (2.10)$$

Letting  $p = 1$  in Cor. 5, we can see

**Corollary 6.** (MacGregor [7]) Let  $f(z), g(z) \in A$  and satisfies

$$\operatorname{Re} \left\{ \frac{zg'(z)}{g(z)} \right\} > 0 \quad (z \in U) . \text{ If } \operatorname{Re} \left\{ \frac{f(z)}{g(z)} \right\} > 0 \quad (z \in U) , \text{ then}$$

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0 \text{ in } |z| < 2 - \sqrt{3} .$$

Now, we prove the next theorem .

**Theorem 5.** Let  $f(z) \in A_p$  and  $a > 0$ . If

$$\operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{z^p} \right\} > 0 \quad (z \in U) ,$$

then

$$\operatorname{Re} \left\{ \frac{L_p(a+2, c)f(z)}{L_p(a+1, c)f(z)} \right\} \geq \frac{(a+1) - 2|z| - (a+1)|z|^2}{(a+1)(1-|z|^2)} . \quad (2.11)$$

**Proof.** We put  $g(z) = \frac{L_p(a+1, c)f(z)}{z^p}$ . Then  $g(z) = 1 + a_1z + a_2z^2 + \dots$  is analytic and satisfies  $\operatorname{Re} \{g(z)\} > 0$  in  $U$ . Differentiating  $z^p \cdot p(z) = L_p(a+1, c)f(z)$  and using Lemma 1, we have

$$p + \frac{zg'(z)}{g(z)} = (a+1) \frac{L_p(a+2, c)f(z)}{L_p(a+1, c)f(z)} - (a+1-p) ,$$

that is,

$$\frac{L_p(a+2, c)f(z)}{L_p(a+1, c)f(z)} = 1 + \frac{1}{a+1} \cdot \frac{zg'(z)}{g(z)} .$$

Therefore, from Lemma 4

$$\begin{aligned} \operatorname{Re} \left\{ \frac{L_p(a+2, c)f(z)}{L_p(a+1, c)f(z)} \right\} &\geq 1 - \frac{1}{a+1} \left| \frac{zg'(z)}{g(z)} \right| \\ &\geq 1 - \frac{1}{a+1} \cdot \frac{2|z|}{1-|z|^2} \\ &= \frac{(a+1) - 2|z| - (a+1)|z|^2}{(a+1)(1-|z|^2)} . \end{aligned}$$

This completes the proof of Thm. 5,

Letting  $a = c = p$  in Thm. 5, we have

**Corollary 7.** Let  $f(z) \in A_p$ . If

$$\operatorname{Re} \left\{ \frac{f'(z)}{z^p - 1} \right\} > 0 \quad (z \in U) ,$$

then

$$1 + \operatorname{Re} \left\{ \frac{zf''(z)}{f'(z)} \right\} > 0 \text{ in } |z| < \frac{\sqrt{1+p^2}-1}{p} . \quad (2.12)$$

**Proof.** By easy calculation, we have

$$\operatorname{Re} \left\{ \frac{L_p(p+2, p)f(z)}{L_p(p+1, p)f(z)} \right\} = \operatorname{Re} \left\{ \frac{2f'(z) + f''(z)}{(p+1)f'(z)} \right\} = \frac{1}{p+1} \left( 2 + \operatorname{Re} \frac{zf''(z)}{f'(z)} \right) .$$

Therefore, from Thm. 5, we have

$$1 + \operatorname{Re} \frac{zf''(z)}{f'(z)} \geq \frac{p - 2|z| - p|z|^2}{1 - |z|^2} \quad (z \in U) .$$

Putting  $p = 1$  in Cor. 7, we have

**Corollary 8.** (MacGregor [6]) Let  $f(z) = z + a_2z^2 + \dots$  be analytic in  $U$ . If

$$\operatorname{Re} \{f'(z)\} > 0 \quad (z \in U) ,$$

then

$$1 + \operatorname{Re} \frac{zf''(z)}{f'(z)} > 0 \text{ in } |z| < \sqrt{2} - 1 .$$

In order to prove next theorem, we need the following lemma due to Ponnusamy .

**Lemma 5.** If  $p(z)$  is analytic in  $U$  with  $p(0) = 1$ , and if  $\lambda$  is a complex number satisfying  $\operatorname{Re} \lambda \geq 0$  ( $\lambda \neq 0$ ), then

$$\operatorname{Re} \{p(z) + \lambda zp'(z)\} > \alpha \quad (0 \geq \alpha < 1) \quad (2.13)$$

implies

$$\operatorname{Re} \{p(z)\} > \alpha + (1 - \alpha)(2\gamma - 1) , \quad (2.14)$$

where  $\gamma$  is given by

$$\gamma = \gamma(\operatorname{Re} \lambda) = \int_0^1 \frac{dt}{1 + t^{\operatorname{Re} \lambda}} . \quad (2.15)$$

is an increasing function of  $\operatorname{Re} \lambda$  and  $\frac{1}{2} \leq \gamma < 1$ . The estimate is sharp in the sense that the bound cannot be improved.

Applying Lemma 5, we prove

**Theorem 6.** Let  $\lambda$  be a complex number satisfying  $\operatorname{Re} \lambda > 0$ . Let  $f(z) \in \mathbf{A}_p$  satisfies the condition

$$\operatorname{Re} \left\{ (1 - \lambda) \left( \frac{L_p(a, c)f(z)}{z^p} \right)^\mu + \lambda \cdot \frac{L_p(a + 1, c)f(z)}{z^p} \left( \frac{L_p(a, c)f(z)}{z^p} \right)^{\mu-1} \right\} > \alpha \quad (z \in U) , \quad (2.16)$$

for some  $\alpha$  ( $0 \leq \alpha < 1$ ) and  $\mu > 0$ . Then

$$\operatorname{Re} \left( \frac{L_p(a, c)f(z)}{z^p} \right)^\mu > \alpha + (1 - \alpha)(2\gamma - 1) \quad (z \in U) , \quad (2.17)$$

where  $\gamma = {}_2F_1 \left( 1, \frac{\mu a}{\operatorname{Re} \lambda}; 1 + \frac{\mu a}{\operatorname{Re} \lambda}; -1 \right)$ . The estimate is sharp.

**Proof.** We put

$p(z) = \left( \frac{L_p(a, c)f(z)}{z^p} \right)^\mu$ , then  $p(z)$  is analytic in  $U$  and  $p(0) = 1$ . Using Lemma 1, we have

$$\begin{aligned} p(z) + \frac{\lambda}{a\mu} zp'(z) &= (1 - \lambda) \left( \frac{L_p(a, c)f(z)}{z^p} \right)^\mu + \lambda \frac{L_p(a + 1, c)f(z)}{z^p} \left( \frac{L_p(a, c)f(z)}{z^p} \right)^{\mu-1} \end{aligned}$$

From assumption,

$$\operatorname{Re} \left\{ p(z) + \frac{\lambda}{a\mu} zp'(z) \right\} > \alpha \quad (0 \leq \alpha < 1, \mu > 0) .$$

Therefore, according to Lemma 5, we have

$$\operatorname{Re} \{p(z)\} > \alpha + (1 - \alpha)(2\gamma - 1) ,$$

where

$$\gamma = \gamma(\operatorname{Re} \frac{\lambda}{a\mu}) = \int_0^1 \frac{dt}{1 + t^{\operatorname{Re}(\lambda/a\mu)}} .$$

We set  $\operatorname{Re} \lambda = \lambda_1$ , then we have

$$\begin{aligned} \gamma &= \int_0^1 \frac{dt}{1 + t^{\lambda_1/a\mu}} = \frac{a\mu}{\lambda_1} \int_0^1 \mu^{a\mu/\lambda_1-1} (1 + \mu)^{-1} du \\ &= {}_2F_1(1, \frac{a\mu}{\lambda_1}; 1 + \frac{a\mu}{\lambda_1}; -1) . \end{aligned}$$

Putting  $\mu = 1, \lambda = 1$  in Thm. 6, we have

**Corollary 9.** Let  $f(z) \in \mathbf{A}_p$  satisfies the condition

$$\operatorname{Re} \left\{ \frac{L_p(a+1, c)f(z)}{z^p} \right\} > \alpha \quad (z \in U) , \quad (2.18)$$

for some  $\alpha$  ( $0 \leq \alpha < 1$ ). Then

$$\operatorname{Re} \left\{ \frac{L_p(a, c)f(z)}{z^p} \right\} > \alpha + (1 - \alpha)(2\gamma - 1) \quad (z \in U) , \quad (2.19)$$

where  $\gamma = {}_2F_1(1, a; a+1; -1)$ . The estimate is sharp.

Taking  $a = c = 1$  and  $p = 1$  in Cor. 9, we can have the following well-known result.

**Corollary 10.** Let  $f(z) = z + a_2 z^2 + \dots$  be analytic in  $U$  satisfying

$$\operatorname{Re} \{f'(z)\} > \alpha \quad (0 \leq \alpha < 1) . \quad (2.20)$$

Then we have

$$\operatorname{Re} \left\{ \frac{f(z)}{z} \right\} > \alpha + (1 - \alpha)(2\gamma - 1) , \quad (2.21)$$

where

$$\gamma = {}_2F_1(1, 1; 2; -1) = \ln 2 .$$

## References

1. S. D. Bernardi, *Convex and starlike univalent functions*, Trans. A. M. S. 135(1969), 429-446.
2. B. C. Carlson and D. B. Shaffer, *Starlike and prestarlike hypergeometric functions*, SIAMJ. Math. Anal. 15(1984), 737-745.
3. P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften 259, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.
4. R. J. Libera, *Some classes of regular univalent functions*, Proc. A. M. S. 16(1965), 755-758.
5. A. E. Livingston, *On the radius of univalence of certain analytic functions*, Proc. A. M. S. 17(1966), 352-357.
6. T. H. MacGregor, *Functions whose derivative has a positive real part*, Trans. Amer. Math. Soc. 104(1962), 532-537.
7. T. H. MacGergor, *The radius of univalence of certain analytic functions I*, Proc. Amer. Math. Soc. 14(1963), 514-520.
8. S. Ruscheweyh, *New criteria for univalent functions*, Proc. A. M. S. 49(1945), 109-115.
9. H. Saitoh, *Some properties of certain multivalent functions*, Tsukuba J. Math. 15(1991), 105-111.
10. H. Saitoh, *On certain class of multivalent functions*, Math. Japon. 37(1992), 871-875.
11. H. Saitoh, *A linear operator and its applications of first order differential subordinations*, to appear in Math. Japonica.
12. H. Saitoh and M. Nunokawa, *On functions satisfying  $\operatorname{Re} \{f(z)/z^p\} > 0$* , Bull. Soc. Royale Sci. Liege, 56(1987), 462-464.
13. H. Saitoh, M. Nunokawa, S. Owa et al, *A note on certain class of analytic functions satisfying  $\operatorname{Re} \{f(z)/z^p\} > 0$* , Math. Today, 7(1989), 33-36.
14. H. Saitoh, S. Owa et al, *An application of a certain integral operator*, Applied Math. Letters, 5(1992), 21-24.
15. H. M. Srivastava and S. Owa, *Some characterization and distortion theorems involving fractional calculus, generalized hypergeometric functions, Hadamard products, linear operators, and certain subclasses of analytic functions*, Nagoya Math. J. 106(1987), 1-28.
16. K. Yamaguchi, *On functions satisfying  $\operatorname{Re} \{f(z)/z\} > 0$* , Proc. Amer. Math. Soc 17(1966), 588-591.